

See discussions, stats, and author profiles for this publication at:
<http://www.researchgate.net/publication/270653925>

Effects of an eight-week whole body vibration on lower extremity muscle tone and function in children with cerebral palsy

ARTICLE *in* RESEARCH IN DEVELOPMENTAL DISABILITIES · JANUARY 2015

Impact Factor: 3.4 · DOI: 10.1016/j.ridd.2014.12.017 · Source: PubMed

5 AUTHORS, INCLUDING:



[Hsin-Yi Kathy Cheng](#)

Chang Gung University

28 PUBLICATIONS 127 CITATIONS

[SEE PROFILE](#)



[Yu-Chun Yu](#)

National Taoyuan Special School, Tao...

3 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



[Yan-Ying Ju](#)

National Taiwan Sport University

19 PUBLICATIONS 107 CITATIONS

[SEE PROFILE](#)



Effects of an eight-week whole body vibration on lower extremity muscle tone and function in children with cerebral palsy



Hsin-Yi Kathy Cheng^a, Yu-Chun Yu^b, Alice May-Kuen Wong^c, Yung-Shen Tsai^d, Yan-Ying Ju^{e,*}

^a Graduate Institute of Early Intervention, College of Medicine, Chang Gung University, Tao-Yuan, Taiwan

^b National Taoyuan Special School, 10 Te-Shou St., Tao-Yuan 330, Taiwan

^c Department of Physical Medicine and Rehabilitation, Chang Gung Memorial Hospital, Tao-Yuan 330, Taiwan

^d Graduate Institute of Sports Equipment Technology, University of Taipei, Taipei, Taiwan

^e Department of Adapted Physical Education, National Taiwan Sport University, 250 Wen-Hua 1st Rd., Kwei-Shan, Tao-Yuan 333, Taiwan

ARTICLE INFO

Article history:

Received 4 November 2014

Accepted 9 December 2014

Available online 7 January 2015

Keywords:

Cerebral palsy

Spasticity

Whole body vibration

Ambulation

Rehabilitation

ABSTRACT

The aim of this study was to evaluate the effect of an eight-week whole body vibration (WBV) on lower extremity spasticity and ambulatory function in children with cerebral palsy with a complete crossover design. Sixteen participants aged 9.2 (2.1) years participated in this study. Half of the participants received a 10-min WBV, 3 times a week for 8 weeks. Then a 4-week washout period followed, after which they received a sham WBV 3 times a week for 8 weeks. The other half received the intervention in a reversed order. The participants were evaluated via variables measuring range-of-motion, muscle tone, and ambulatory function before, immediately after, 1 day after, and 3 days after each intervention. Repeated-measures analyses revealed significant beneficial effects on most variables except the passive range-of-motion measurement. Significant correlations were found between timed up-and-go and relaxation index, and between timed up-and-go and six-minute walk test. The results suggested that an 8-week WBV intervention normalized muscle tone, improved active joint range and enhanced ambulatory performance in children with cerebral palsy for at least 3 days. These indicated that regular WBV can serve as an alternative, safe, and efficient treatment for these children in both clinical and home settings.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Cerebral palsy (CP) is the most common childhood disabilities which affect individual's posture and movement (Koman, Smith, & Shilt, 2004). Compared to the typically developed children, these children have impaired sensation and increased muscle tone therefore they have trouble voluntarily controlling their muscles. About seventy to eighty percent of children with CP demonstrate spastic clinical features (Kriger, 2006). Traditionally spasticity is managed via anti-spastic medication

* Corresponding author. Tel.: +886 3 328 3201x8624; fax: +886 3 327 7475.

E-mail addresses: kcheng@mail.cgu.edu.tw (H.-Y. Cheng), ot.spring.yu@gmail.com (Y.-C. Yu), walice@adm.cgmh.org.tw (A.-K. Wong), ytsai@utapei.edu.tw (Y.-S. Tsai), yanju@ntsu.edu.tw (Y.-Y. Ju).

or injection. Several disadvantages are associated with these treatments. First, not all patients achieve satisfactory results, and not all parents are comfortable with long-term medication use or invasive injection. In addition, there are side effects such as general weakness that in turn affect their functional performance. As a result, there is a need for alternative or additional treatment options.

Other than spasticity, poor muscle strength is also an important factor that leads to movement dysfunction in CP (Damiano, Martellotta, Sullivan, Granata, & Abel, 2000). The weak quadriceps in this population inevitably impaired their walking performance (Damiano, Kelly, & Vaughn, 1995). These days whole body vibration (WBV) has become popular in local gym and fitness centers, and it was found to offer fitness and health benefits, including flexibility and muscle strength (Dolny & Reyes, 2008; Kosar, Candow, & Putland, 2012; Rittweger, 2010). Recently, it seems to reach satisfactory results in managing spasticity and muscle strength among participants with upper motor neuron syndrome (Ahlborg, Andersson, & Julin, 2006; Chan et al., 2012; Ness & Field-Fote, 2009).

Whole body vibration (WBV) is a training method which exposes the whole body of an individual to low frequency, low amplitude mechanical stimuli via a vibrating platform. The vibration stimulates the muscle spindles, sending nerve impulses to initiate muscle contractions according to the tonic vibration reflex (Cardinale & Bosco, 2003). Its effects on enhancing health and fitness in general population have been studied extensively; however, little was done with special groups, for example children with CP. Ruck, Chabot, and Rauch (2010) examined the effects of a 9-min WBV program on children with CP and found their mobility improved (Ruck et al., 2010). In this particular study, the only indicator for mobility was the change in self-selected walking speed (Ruck et al., 2010). Unger, Jelsma, and Stark (2013) investigated the vibration on trunk muscle strengthening and found beneficial effects onto posture and gait. Other than the postural related measurements, there was only one gait parameter, the 1-min walk test, assessed (Unger et al., 2013). Furthermore, the direct effect on spasticity was never evaluated.

Ahlborg et al. (2006) performed a comprehensive survey of spastic, strength, and walking variables after an 8-week WBV or an 8-week resistance training in adult with CP (Ahlborg et al., 2006). They found WBV was more effective in decreasing spasticity of the knee extensors compared to the resistance training. Leg muscle strength increased in both groups. On the other hand, the ambulatory parameters did not change significantly. Although this study demonstrated an extensive investigation on the effects of WBV on CP, there were aspects that can be improved. First, findings of the study were gathered from adults with CP, not children. Children usually demonstrate a better potential in functional improvement, therefore studies with children as subjects are of great needs. Second, this study had no control group. A control group provides true baselines for all testing variables, and this becomes substantially important when dealing with children. Third, when isokinetic strength was measured before ambulatory parameters, the muscle exertion would hinder the effects of WBV on ambulation. Last, the study only measured immediate influences, no lasting effects were evaluated.

Therefore, the aim of this study was to evaluate the immediate and lasting effects of WBV on lower extremity spasticity and ambulatory function in children with CP. It was hypothesized that upon the completion of this 8-week program, the spasticity among these CP children would decrease significantly and their ambulatory function would improve.

2. Materials and methods

A crossover repeated measures design was employed in this study. Sixteen children with CP were randomly divided into two groups. One group received an 8-week WMV intervention followed by an 8-week control condition, with a 4-week rest in between; the other group began their treatment sequence with the control condition to counterbalance the order effects (Fig. 1). The outcome variables included active and passive range-of-motion (AROM and PROM) of both knees, relaxation




Fig. 1. Flow chart of the crossover design.

index (RI) assessed by the Wartenburg Pendulum test, modified Ashworth scales (MAS) for bilateral knee extensors, timed up-and-go (TUG) and 6-min walk test (6WMT).

2.1. Participants

Sixteen children with CP, eight boys and eight girls, aged 9.2 (2.1) years with diagnoses of either spastic diplegia ($N = 11$) or spastic quadriplegia ($N = 5$) were recruited from a local hospital and a special education school. The inclusion criteria were: (1) add-up MAS scores of bilateral lower extremity greater than two; (2) ability to walk for at least 6 min with or without walking aids; and (3) ability to comprehend commands. Criteria which would affect their muscle tone and ambulatory performance were excluded. The exclusion criteria were: (1) the presence of any progressive disorder or a severe concurrent illness not typically associated with CP; (2) lower extremity surgery within the past 6 months; (3) nerve block or botulinum toxin injection within the past 3 months; (4) a limitation of knee joint range-of-motion over 10° ; (5) epilepsy. One of the authors performed the selection process. Most participants walked independently without walking aids except four, who used 4-wheeled walkers for ambulation. The Institutional Review Board for Human Studies of Chang Gung Memorial Hospital approved this protocol. Written informed consents were obtained from all participants and their legal guardians.

2.2. Equipment

The equipment utilized in this study included electrogoniometers (SG110, Biometrics, Ltd, Cwmfelinfach, Gwent, UK) and a WBV platform (AV-001A, Body Green, Taipei, Taiwan). The electrogoniometers, taped on the lateral sides of the knee joints, were used to measure the degree of flexion and extension. A data acquisition unit (MP150, Biopac Systems Inc., Santa Barbara, CA, USA) was used for signal registration. The WBV platform could provide vertical vibrations with frequencies and amplitudes ranging from 8 to 40 Hz and 0.4 to 2.0 mm, respectively.

2.3. Protocol

The vibration platform located in a quiet room with the environment set up for variable measurements. Subject was transferred to the room via wheelchair to avoid any fatigue prior to the testing. All participants underwent a series of outcome measurements in the order of AROM and PROM for knee joints, Pendulum test, MAS, TUG, 6WMT. Variables measured before the intervention (Time1) were the baseline scores. These measurements were administered again immediately after 8-week intervention (Time2), 1 day after (Time3), and 3 days after (Time4) the 8-week period. The test-retest reliability for each variable was evaluated from two consecutive days before intervention with ten children with spastic cerebral palsy, aged 9.7 (1.7) years. Change scores (Diff2-1/Diff3-1/Diff4-1) were used to show the improvement of each dependent variable measured in time 2, 3 and 4 compared to the baseline. The duration of each assessment was about 45 min.

Each subject performed both the WBV and control conditions in a counterbalanced order on two separate days, one week apart. The subject was instructed to adhere to the daily routines. Prior to the first experimental trial, the participant was requested to stand on the vibration platform with the knees slightly flexed at about 30 degrees from full extension. Anterior knee stops and a pelvic belt were used along with the WBV machine for securing their knees and pelvis to provide external support while standing. The participants were instructed to avoid holding on to the supported rail if possible however they were allowed to hold onto the rail if necessary (Ness & Field-Fote, 2009). They were asked to focus on standing with equal weights on both legs, and keep body movements to a minimum. Passive vibration was then delivered at 20 Hz with a vertical displacement of 2 mm. The WBV was given for 10 min. For the control condition, participants followed the same procedure instead that the vibration machine was not turned on. After the intervention, the participants were moved to the assessment area in wheelchairs to avoid any physical activity on their feet.

2.4. Outcome variables

2.4.1. Measurement of joint range-AROM & PROM

The AROM measures the range of movement through which the subject can actively move a joint using the adjacent muscles, whereas the PROM measures the amount of motion at a given joint when the joint is moved by an external force. Electrogoniometers were secured on the lateral sides of the knees for signal registration. For the AROM measurement, the subject was first asked to actively move the knee to full extension and then full flexion. Subject was asked to maintain full extension and flexion for 3 s each and the median angles of the two 3-s intervals were used for AROM calculation. For the PROM condition, their knees were passively moved by an examiner to the end range. The intraclass correlation coefficients (ICCs) for AROM and PROM were 0.73 and 0.79, respectively.

2.4.2. Measurement of spasticity—Wartenburg Pendulum test

The Pendulum test evaluated muscle tone by using gravity to provoke stretch reflexes of knee extensors during passive swinging of the lower limb. The oscillatory movements of the lower leg were captured by electrogoniometer at the lateral knee. The relaxation index (RI) was then calculated as follows: $RI = (\text{starting angle} - \text{first angle}) / (\text{starting angle} - \text{resting angle})$

angle) \times 1.6 (BurrIDGE et al., 2005). The ICC calculated for RI was 0.72. In case of spasticity, the oscillatory amplitudes were significantly less as compared to those with normal muscle tone.

2.4.3. Measurement of spasticity-MAS

The MAS scale is widely used method for measuring spasticity. A score of 0–4 was used to assess the resistance of knee extensors to passive movement. The children lay supine with the lower leg hanging freely over the edge and the knee passively extended by the examiner. The examiner moved the subject's knee to a position of maximal flexion over 1 s. A score was then given based on the classification by Bohannon and Smith (Bohannon & Smith, 1987). The MAS scores of both legs were added up to represent the muscle tone of the subject's lower extremities. The ICC calculated for MAS was 0.92.

2.4.4. Functional measurement-TUG

TUG is a test used to assess a person's mobility. TUG measured the time required for an individual to stand up from a chair with armrests, walk 3 m, turn, walk back to the chair, and sit down. The subject walked with their regular footwear at their preferred speed with or without walking aids. The examiner stayed with the participants to protect their safety at all times. High test–retest reliability (ICC = 0.94) and adequate validity (spearman $r = -0.77$) for the TUG in children with CP have been reported (Zaino, Marchese, & Westcott, 2004).

2.4.5. Functional measurement-6MWT

Six-MWT measured the distance walked within 6 min. A 7-m walkway was marked on a level floor. The subject was asked to walk continuously along this walk way, turn around at the end, and continue to cover as much ground as possible over 6 min. The examiner spoke standardized phrases to the patient to avoid improper encouragement and enthusiasm. High test–retest reliability (ICC = 0.98) for the 6MWT in children with CP was revealed (Maher, Williams, & Olds, 2008).

2.5. Statistics

Data were analyzed using the statistical analysis software SPSS (version 17.0, SPSS Inc., Chicago, IL, USA). Descriptive statistics were used to calculate participants' demographics. A two-way repeated measures analysis of variance (ANOVA) (condition(2) \times change scores of time (3)) was performed for all variables. The alpha level was set at 0.05. Multiple comparisons between pairs of means were used to determine where the differences existed with Bonferroni adjustments of alpha level. The relationship between functional improvements and changes in spasticity were examined by correlation analysis.

3. Results

The initial spasticity measurements for MAS and RI were 4.61 (0.95) and 0.59 (0.07), respectively. The initial measurements for TUG and 6MWT were 13.53 (2.30) s and 201.33 (17.51) m, respectively.

Descriptive and inferential statistics for outcome analyses are presented in Table 1. Significant differences were found in change scores between the treatment and control condition for variables including MAS, RI, and 6MWT. Multiple comparisons revealed that differences existed between change scores Diff2-1 and Diff4-1 only for MAS ($p = 0.036$) and RI ($p = 0.022$); and between change scores Diff2-1 and Diff3-1 ($p = 0.006$), Diff2-1 and Diff4-1 ($p = 0.012$), but not Diff3-1 and Diff4-1 ($p = 0.907$), for 6MWT. For AROM and TUG, significant differences were found in condition only. However, no difference was found in PROM.

Correlation of scores was performed to reveal the relationship between functional improvement and changes in muscle tone. Correlation results revealed that change in TUG was significantly correlated with change in RI ($r = -0.526$, $p = 0.036$),

Table 1
Descriptive and inferential statistics for outcome analyses.

Variables	WBV condition			Control condition			$P_{\text{condition}}$	$P_{\text{change_score}}$
	Change_score ^a			Change_score ^a				
	Diff2-1	Diff3-1	Diff4-1	Diff2-1	Diff3-1	Diff4-1		
AROM (°)	5.74 (3.46)	4.35 (2.05)	3.34 (2.73)	0.24 (2.29)	0.61 (2.39)	0.22 (2.13)	0.000**	0.154
PROM(°)	1.13 (2.04)	-0.75 (2.88)	0.48 (2.02)	0.25 (1.23)	0.09 (1.23)	0.33 (1.26)	0.892	0.118
RI	0.13 (0.06)	0.09 (0.06)	0.09 (0.06)	0.00 (0.04)	0.02 (0.04)	-0.00 (0.30)	0.000**	0.030*
MAS	-1.43 (1.05)	-1.20 (1.11)	-0.50 (0.73)	-0.17 (0.25)	-0.13 (0.34)	0.03 (0.46)	0.000**	0.033*
TUG (s)	-2.49 (1.56)	-2.79 (1.68)	-2.39 (1.52)	-0.39 (1.28)	-0.52 (1.37)	-0.54 (1.18)	0.000**	0.416
6MWT (m)	16.89 (8.01)	6.96 (6.74)	7.66 (6.46)	-0.20 (7.50)	-1.70 (9.31)	0.57 (4.25)	0.000**	0.005**

Values are expressed as mean \pm SD.

^a Different score Diff2-1: difference between measurements "Time2" & "Time1". Diff3-1: difference between measurements "Time3" & "Time1". Diff4-1: difference between measurements "Time4" & "Time1".

* $p < 0.05$.

** $p < 0.01$.

but not with change in MAS ($r = -0.129$, $p = 0.633$). Six-minute walk test was significantly correlated with change in TUG only ($r = -0.713$, $p = 0.002$), but not with changes in RI or MAS. No other correlations were significant.

4. Discussion

This study is the first to implement a crossover repeated measures design for measuring the effects of an 8-week WBV protocol on children with CP. The results were consistent with our hypothesis that spasticity (MAS and RI) would reduce, and ambulatory function (TUG and 6MWT) would improve significantly after the WBV intervention. These effects could last up to 3 days after the intervention. These benefits would facilitate children to actively engage in daily activities, exercise trainings and therapeutic interventions.

The benefits of the 8-week WBV program are manifold. First, it can suppress spasticity in children with CP. This finding was in line with previous research (Ahlborg et al., 2006; Chan et al., 2012; Liepert and Binder, 2010; Miyara et al., 2014; Ness & Field-Fote, 2009). Using an electrophysiological approach, Liepert and Binder (2010) revealed the vibration effects on target muscles (Liepert and Binder, 2010). Vibration effects are mediated through the activation of muscle spindles and transmission by Ia fibers, which enhances cortical excitability of the vibrated muscle. The vibration stimuli simultaneously reduce activity in antagonistic muscles via reciprocal inhibition and supraspinal inhibition, therefore reducing antagonistic muscle hyperactivity. A more balanced interaction between flexors and extensors can then be achieved. In addition, WBV can induce presynaptic inhibition. Previous studies demonstrated that a vibratory stimulus applied directly to a muscle will induce the presynaptic inhibition of Ia afferents (Schieppati, 1987) and is likely to reduce the release of neurotransmitters to the motoneurons (Katz, 1999), thereby decreasing the monosynaptic reflex excitability.

Second, it significantly improved the AROM in the knees. The vibration delivered directly to the soles of feet, and went all the way up to the subject's body, especially to the lower extremities. These mechanical vibrations stimulated skin receptors and muscle spindles, enhancing the cortical excitability of the vibrated muscle and therefore increasing joint AROM (Katusic, Alimovic, & Mejaski-Bosnjak, 2013). On the other hand, no such improvements occurred in PROM. Clinically, measurement of PROM is assessed by passively moving a joint to its extreme position. This movement is slow and should not elicit reflex activity. This indicated that the value of PROM cannot be interfered with spasticity. Rather, the limitation of PROM is mainly originated from the mechanical constraints, such as the length of musculature surrounding the joint, the flexibility of joint capsule, and the contour of bones.

Third, it can improve walking performance. TUG was significantly improved immediately after and 30-min after the WBV intervention. The improvement on 6MWT was also significant but it attenuated over time after the cease of the intervention. Previously a correlation was found between muscle tone, functional gait parameters and a more normal electromyography pattern (Hesse et al., 1996), indicating that a reduction in spasticity might lead to improvements in motor function and walking ability. Another study also found patients with measurable spasticity showed reduced knee angular velocity during walking since their muscles were rather stiff (Tuzson, Granata, & Abel, 2003). As a consequence, the ambulation function improved as the spasticity decreased. In addition, results from the current study also revealed significant correlation between change in TUG and change in RI. The improvement in RI remained during follow-up period of 30 min, which paralleled with the improvement in TUG. Unlike TUG, 6MWT might require better endurance which could not be acquired from this intervention. Thus the improvement on 6MWT might fade faster in this protocol.

Vibrations are believed to initiate muscle contractions by stimulating muscle spindles and alpha motor neurons, resulting in an effect similar to that of conventional resistance training (Delecluse, Roelants, & Verschueren, 2003). In the current study the WBV was performed on a vibrating platform during supported standing. It is a form of muscle and proprioceptive training that is largely independent of the motivation of the participant (Rauch, 2009). Children with young ages are usually less motivated since they do not understand well the link between therapeutic training and health benefits. Therefore WBV can serve as a substitute for traditional resistance training and rehabilitation programs.

Based on the subjective reports of the participants, negative side-effects seldom occur with the WBV intervention. One participant reported stiff legs and one reported back muscle soreness in the first two sessions of WBV intervention. Still, attentions need to be paid. Vibrating platform might threaten participants' balance and cause their feet to slide off. Close supervision is needed especially for those with physical/sensory disabilities and those with young ages.

There are still limitations in this study. First, results from this study revealed the effects of WBV intervention in decreasing spasticity and improving walking function, but whether the effects of WBV are comparable to those provided by oral medication or muscle injection are still questionable. Future studies should directly compare the antispastic effects of vibration to those of antispastic agents. Second, participants in this study did not cease their regular therapeutic activities. The WBV program was added to their daily routines. The authors controlled the possible influences via crossover design. Even that, the effects from their physical exercises or school activities cannot be completely removed.

5. Conclusion

To sum up, WBV program can be a suitable approach to control spasticity and improve walking function in children with CP, particularly it is non-invasive, easy to apply, relatively safe and requires little efforts. It may also serve as a routine addition or substitute for other anti-spastic agents with future supporting studies. Performed before the rehabilitation session, it serves as a spasticity reducing protocol and gets the individual ready for advanced exercise trainings. Previous

studies also revealed other benefits of WBV on various physiological measures, including the oxygen consumption, muscle temperature, skin blood flow, muscle power, muscle strength, balance, and bone density (Cochrane, Stannard, Sargeant, & Rittweger, 2008; Lohman Iii, Petrofsky, Maloney-Hinds, Betts-Schwab, & Thorpe, 2007; Rauch, 2009). These together support the potential clinical application of WBV in CP rehabilitation.

Acknowledgments

The authors thank all children, parents, school teachers and staff for their participation and assistance in this study. This study was supported by the Ministry of Science and Technology (103-2221-E-182-048, 103-2410-H-179-002), and the Chang Gung Medical Research Program, Taiwan (CMRPD1D0131).

References

- Ahlborg, L., Andersson, C., & Julin, P. (2006). Whole-body vibration training compared with resistance training: Effect on spasticity, muscle strength and motor performance in adults with cerebral palsy. *Journal of Rehabilitation Medicine*, 38(5), 302–308. <http://dx.doi.org/10.1080/16501970600680262>
- Bohannon, R. W., & Smith, M. B. (1987). Assessment of strength deficits in eight paretic upper extremity muscle groups of stroke patients with hemiplegia. *Physical Therapy*, 67(4), 522–525.
- Burridge, J. H., Wood, D. E., Hermens, H. J., Voerman, G. E., Johnson, G. R., van Wijck, F., et al. (2005). Theoretical and methodological considerations in the measurement of spasticity. *Disability and Rehabilitation*, 27(1–2), 69–80.
- Cardinale, M., & Bosco, C. (2003). The use of vibration as an exercise intervention. *Exercise and Sport Sciences Reviews*, 31(1), 3–7.
- Chan, K. S., Liu, C. W., Chen, T. W., Weng, M. C., Huang, M. H., & Chen, C. H. (2012). Effects of a single session of whole body vibration on ankle plantarflexion spasticity and gait performance in patients with chronic stroke: A randomized controlled trial. *Clinical Rehabilitation*, 26(12), 1087–1095.
- Cochrane, D. J., Stannard, S. R., Sargeant, A. J., & Rittweger, J. (2008). The rate of muscle temperature increase during acute whole-body vibration exercise. *European Journal of Applied Physiology*, 103(4), 441–448.
- Damiano, D. L., Martellotta, T. L., Sullivan, D. J., Granata, K. P., & Abel, M. F. (2000). Muscle force production and functional performance in spastic cerebral palsy: Relationship of cocontraction. *Archives of Physical Medicine and Rehabilitation*, 81(7), 895–900.
- Damiano, D. L., Kelly, L. E., & Vaughn, C. L. (1995). Effects of quadriceps femoris muscle strengthening on crouch gait in children with spastic diplegia. *Physical Therapy*, 75(8), 658–667.
- Delecluse, C., Roelants, M., & Verschueren, S. (2003). Strength increase after whole-body vibration compared with resistance training. *Medicine and Science in Sports and Exercise*, 35(6), 1033–1041.
- Dolny, D. G., & Reyes, G. F. C. (2008). Whole body vibration exercise: Training and benefits. *Current Sports Medicine Reports*, 7(3), 152–157.
- Hesse, S., Krajnik, J., Luecke, D., Jahnke, M. T., Gregoric, M., & Mauritz, K. H. (1996). Ankle muscle activity before and after botulinum toxin therapy for lower limb extensor spasticity in chronic hemiparetic patients. *Stroke*, 27(3), 455–460.
- Katusic, A., Alimovic, S., & Mejaski-Bosnjak, V. (2013). The effect of vibration therapy on spasticity and motor function in children with cerebral palsy: A randomized controlled trial. *NeuroRehabilitation*, 32(1), 1–8. <http://dx.doi.org/10.3233/NRE-130817>
- Katz, R. (1999). Presynaptic inhibition in humans: A comparison between normal and spastic patients. *Journal of Physiology-Paris*, 93(4), 379–385.
- Koman, L. A., Smith, B. P., & Shilt, J. S. (2004). Cerebral Palsy. *Lancet*, 363, 1619–1631.
- Kosar, A. C., Candow, D. G., & Putland, J. T. (2012). Potential beneficial effects of whole-body vibration for muscle recovery after exercise. *The Journal of Strength & Conditioning Research*, 26(10), 2907–2911.
- Krigger, K. W. (2006). Cerebral palsy: An overview. *American Family Physician*, 73(1), 91–100.
- Liepert, J., & Binder, C. (2010). Vibration-induced effects in stroke patients with spastic hemiparesis—A pilot study. *Restorative Neurology and Neuroscience*, 28, 729–735.
- Lohman Iii, E. B., Petrofsky, J. S., Maloney-Hinds, C., Betts-Schwab, H., & Thorpe, D. (2007). The effect of whole body vibration on lower extremity skin blood flow in normal subjects. *Medical Science Monitor*, 13(2), CR71–CR76.
- Maher, C. A., Williams, M. T., & Olds, T. S. (2008). The six-minute walk test for children with cerebral palsy. *International Journal of Rehabilitation Research*, 31(2), 185–188.
- Miyara, K., Matsumoto, S., Uema, T., Hirokawa, T., Noma, T., Shimodozono, M., et al. (2014). Feasibility of using whole body vibration as a means for controlling spasticity in post-stroke patients: A pilot study. *Complementary Therapies in Clinical Practice*, 20(1), 70–73.
- Ness, L. L., & Field-Fote, E. C. (2009). Effect of whole-body vibration on quadriceps spasticity in individuals with spastic hypertonia due to spinal cord injury. *Restorative Neurology and Neuroscience*, 27(6), 621–631.
- Rauch, F. (2009). Vibration therapy. *Developmental Medicine and Child Neurology*, 51(Suppl. 4), 166–168. <http://dx.doi.org/10.1111/j.1469-8749.2009.03418.x>
- Rittweger, J. (2010). Vibration as an exercise modality: How it may work, and what its potential might be. *European Journal of Applied Physiology*, 108(5), 877–904.
- Ruck, J., Chabot, G., & Rauch, F. (2010). Vibration treatment in cerebral palsy: A randomized controlled pilot study. *Journal of Musculoskeletal Neuronal Interactions*, 10(1), 77–83.
- Schieppati, M. (1987). The Hoffmann reflex: A means of assessing spinal reflex excitability and its descending control in man. *Progress in Neurobiology*, 28(4), 345–376.
- Tuzson, A. E., Granata, K. P., & Abel, M. F. (2003). Spastic velocity threshold constrains functional performance in cerebral palsy. *Archives of Physical Medicine and Rehabilitation*, 84(9), 1363–1368.
- Unger, M., Jelsma, J., & Stark, C. (2013). Effect of a trunk-targeted intervention using vibration on posture and gait in children with spastic type cerebral palsy: A randomized control trial. *Developmental Neurorehabilitation*, 16(2), 79–88. <http://dx.doi.org/10.3109/17518423.2012.715313>
- Zaino, C. A., Marchese, V. G., & Westcott, S. L. (2004). Timed up and down stairs test: Preliminary reliability and validity of a new measure of functional mobility. *Pediatric Physical Therapy*, 16(2), 90–98.